

# Nutrient Needs at a Glance

Extension Nutrition Specialists  
 The Texas A&M System



## Glossary

<b>Adequate Intake (AI):</b>	set when there is no data to set the RDA	<b>Hemorrhagic:</b>	loss of blood from blood vessels
<b>Acceptable Macronutrient Distribution Range (AMDR):</b>	range of intake for an energy source that reduces risk of chronic disease while providing essential nutrients. Excess leads to weight gain and increased risk of chronic disease.	<b>Ketosis:</b>	a condition caused by abnormal burning of fat in the body
<b>Anorexia:</b>	loss of appetite	<b>Macronutrients:</b>	nutrients—proteins, fats, carbohydrates, others—needed by the body in large amounts
<b>Antioxidant:</b>	a substance that prevents the deterioration or rancidity of fats	<b>Microgram (µg - mcg):</b>	one millionth of a gram
<b>Ataxia:</b>	inability to coordinate voluntary muscles	<b>Milligram (mg):</b>	one thousandth of a gram
<b>Cachexia:</b>	general physical wasting and malnutrition	<b>Neural Tube Defects (NTD):</b>	birth defects due to failure of the neural tube to develop properly during fetal development
<b>Cheilosis:</b>	cracks at the corner of the mouth	<b>Osteomalacia:</b>	softening of bones in adults
<b>Coenzyme:</b>	compound that forms the actual part in an enzyme after combining with a protein component	<b>Osteoporosis:</b>	porous, brittle bones
<b>Daily Values (DVs):</b>	the amount of a nutrient needed daily as determined by the Food and Drug Administration (FDA)	<b>Photophobia:</b>	sensitivity to light
<b>Dermatitis:</b>	inflammation of the skin	<b>Recommended Dietary Allowances (RDA):</b>	the amount of nutrients needed to promote good growth and optimum health in people ages 25 to 50
<b>Desquamation:</b>	loss of a layer of skin	<b>Rickets:</b>	bone deformation in children
<b>Dietary Reference Intakes (DRIs)</b>	general term for a set of reference values for planning and assessing nutrient intakes of healthy people	<b>Scurvy:</b>	weakened cartilages and connective tissue
<b>Eczema:</b>	an inflammatory condition of the skin characterized by redness and itching	<b>Tolerable Upper Intake Level (UL):</b>	highest daily intake that will not cause adverse effects
<b>Edema:</b>	abnormal accumulation of fluid in the body	<b>Xerophthalmia:</b>	an eye condition that can lead to blindness
<b>Glucose Tolerance Factor (GTF):</b>	a dietary agent that facilitates the reaction of insulin		
<b>Gram (g):</b>	metric unit of mass equal to one thousandth (10 <sup>-3</sup> ) of a kilogram		

## References

Data compiled by the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes for Nutrients Reports ([www.nap.edu](http://www.nap.edu)), the Food and Nutrition Board, Institute of Medicine, National Academy of Sciences, Washington, DC: National Academy Press, 1997-2010.

Center for Nutrition Policy and Promotion ([www.cnpp.usda.gov/dietaryguidelines.htm](http://www.cnpp.usda.gov/dietaryguidelines.htm)); Office of Dietary Supplements, National Institute of Health, 2010.

Revised by Mary Kinney Bielamowicz, PhD, RD, LD, Regents Fellow, Professor and Extension Nutrition Specialist, and Sharon F. Robinson, PhD, RD, LD, Associate Professor and Extension Nutrition Specialist, The Texas A&M System, assisted by Dietetic Interns Kelsey Kinsella, Misty Cram, Kelly Vaughan, 2009; Molly Cernosek, 2011.

## Estimated safe and adequate daily dietary intakes of selected vitamins and minerals

DRI's	Age range	RDA* (bold)/AI*		AMDR*	Functions in the body	Sources	Deficiency
Nutrients (macro*)		Males	Females	M + F			
<b>Protein</b> (g/d) (grams*/day)	1–8 years	<b>13–19</b>	<b>13–19</b>	5–30	<ul style="list-style-type: none"> <li>Builds and repairs all body tissue</li> <li>Helps build blood</li> <li>Helps form antibodies to fight infection</li> <li>Supplies food energy at 4 calories per gram</li> </ul>	Animal protein: meat, fish, poultry, eggs, milk, cheese, yogurt Vegetable protein: legumes (peas, beans), whole grain breads and cereals, nuts, peanut butter, soy	Fatigue, loss of appetite, edema*, poor growth
	9–18 years	<b>34–52</b>	<b>34–46</b>	10–30			
	19–50 years	<b>56</b>	<b>46</b>	10–35			
	51–70 years	<b>56</b>	<b>46</b>	10–35			
<b>Fat</b> (g/d)	1–8 years	–	–	25–40	<ul style="list-style-type: none"> <li>Supplies 9 calories per gram (more energy in a small amount of food)</li> <li>Transports fat-soluble vitamins and essential fatty acids needed for body's proper use and storage of fat</li> </ul>	Butter, margarine, shortening, oil, salad dressing, palm and coconut oil, egg yolk, meat with fat, whole milk, cheese, peanut butter	Eczema*, retarded growth, diarrhea, loss of hair
	9–18 years	–	–	25–35			
	19–50 years	–	–	25–35			
	51–70 years	–	–	20–35			
<b>Carbohydrates</b> (g/d)	1–8 years	<b>130**</b>	<b>130**</b>	45–65	<ul style="list-style-type: none"> <li>Supply energy at 4 calories per gram to all body cells</li> <li>Supply glucose to spare protein</li> <li>Help the body use other nutrients</li> </ul>	Breads, cereals, flours, cornmeal, rice, macaroni, noodles, spaghetti, Irish and sweet potatoes, corn, dried fruits, bananas, sugar, syrup, jam, jellies, preserves, honey	Loss of energy, fatigue, ketosis*
	9–18 years	<b>130**</b>	<b>130**</b>	45–65			
	19–50 years	<b>130**</b>	<b>130**</b>	45–65			
	51–70 years	<b>130**</b>	<b>130**</b>	45–65			
<b>Fiber</b> (g/d)	1–8 years	14–20	14–17	None determined	<ul style="list-style-type: none"> <li>May help lower cholesterol</li> <li>Improves bowel motility</li> <li>Gives feeling of fullness without extra calories, promoting satiety and weight loss</li> <li>Contains phytic acids that tie up minerals, which can prevent absorption</li> </ul>	Whole grains (wheat, unmilled rice, oats) or enriched products: cereals, bread, noodles, tortillas, brown rice, oatmeal Vegetables: broccoli, spinach, carrots, beans, peas	Diarrhea; excess fiber makes bulk, which may prevent eating enough food energy or nutrients; high-fiber diets for elderly, very young or those on low-calorie diets may cause nutrient deficiencies
	9–18 years	25–31	22–25				
	19–50 years	31–34	25–28				
	51–70 years	28	22				
Water-soluble vitamins		RDA*/AI*		UL*	Functions in the body	Sources	Deficiency
		Males	Females	M + F			
<b>Vitamin C</b> <b>Ascorbic Acid</b> (mg/d) (milligrams*/day)	1–8 years	<b>15–25</b>	<b>15–25</b>	400–650	<ul style="list-style-type: none"> <li>Helps wounds heal</li> <li>Promotes iron absorption</li> <li>Helps the body maintain collagen (fibrous part of protein for cell structure)</li> <li>Acts as an antioxidant</li> </ul>	All citrus fruits, fruit juices, strawberries, cantaloupe; green or red peppers, raw cabbage, spinach, broccoli, turnip greens, collards, mustard greens, kale, tomatoes, Irish or sweet potatoes	Scurvy*, sore or bleeding gums, poor wound healing, pain in joints, bones, muscles
	9–18 years	<b>45–75</b>	<b>45–65</b>	1,200–1,800			
	19–50 years	<b>90</b>	<b>75</b>	2,000			
	51–70 years	<b>90</b>	<b>75</b>	2,000			
<b>Vitamin B<sub>1</sub> – Thiamin</b> (mg/d)	1–8 years	<b>0.5–0.6</b>	<b>0.5–0.6</b>	None determined	<ul style="list-style-type: none"> <li>Helps the body use carbohydrates for energy</li> <li>Maintains appetite and muscle tone</li> <li>Involved in nervous system function</li> </ul>	Meat (especially pork), liver, heart, kidney, poultry, eggs, milk, dried peas and beans, nuts, whole-grain or enriched bread and cereals	Poor appetite, constipation, depression, apathy, cachexia*, edema*, cardiac failure, cheilosis*
	9–18 years	<b>0.9–1.2</b>	<b>0.9–1.0</b>				
	19–50 years	<b>1.2</b>	<b>1.1</b>				
	51–70 years	<b>1.2</b>	<b>1.1</b>				
<b>Vitamin B<sub>2</sub> – Riboflavin</b> (mg/d)	1–8 years	<b>0.5–0.6</b>	<b>0.5–0.6</b>	None determined	<ul style="list-style-type: none"> <li>Functions as a part of a coenzyme* that assists in energy release</li> <li>Helps in metabolism of amino acids</li> </ul>	Milk, cheese, ice cream, organ meats, eggs, fish, dark green leafy vegetables, enriched breads and cereals	Cheilosis*, scaly desquamation* around nose and ears, sore tongue and mouth, burning and itching eyes, photophobia*
	9–18 years	<b>0.9–1.3</b>	<b>0.9–1.0</b>				
	19–50 years	<b>1.3</b>	<b>1.1</b>				
	51–70 years	<b>1.3</b>	<b>1.1</b>				
<b>Niacin</b> (mg/d NE*) <i>Nicotinic acid</i> <i>Nicotinamide</i>	1–8 years	<b>6–8</b>	<b>6–8</b>	10–15	<ul style="list-style-type: none"> <li>Coenzyme* for carbohydrate metabolism</li> <li>Promotes normal appetite</li> </ul>	Meat, liver, poultry, fish, dried peas and beans, nuts (especially peanuts), whole-grain or enriched cereals and breads, milk, cheese, yogurt	Anorexia*, diarrhea, dermatitis*, confusion, anxiety
	9–18 years	<b>12–16</b>	<b>12–14</b>	20–30			
	19–50 years	<b>16</b>	<b>14</b>	35			
	51–70 years	<b>16</b>	<b>14</b>	35			
<b>Vitamin B<sub>6</sub></b> (mg/d) <i>Pyridoxine</i> <i>Pyridoxal</i> <i>Pyridoxamine</i>	1–8 years	<b>0.5–0.6</b>	<b>0.5–0.6</b>	30–40	<ul style="list-style-type: none"> <li>Coenzyme* for protein utilization</li> <li>Helps convert the amino acid tryptophan to the vitamin Niacin</li> <li>Helps convert complex carbohydrates to simple carbohydrates</li> </ul>	Meat, poultry, fish, sweet potatoes, vegetables, whole grains, fortified cereals	Anemia, nervous irritability, convulsions, weakness, ataxia*, abdominal pain, dermatitis*
	9–18 years	<b>1.0–1.3</b>	<b>1.0–1.2</b>	60–80			
	19–50 years	<b>1.3</b>	<b>1.3</b>	100			
	51–70 years	<b>1.7</b>	<b>1.5</b>	100			
<b>Choline</b> (mg/d)	1–8 years	200–250	200–250	1,000	<ul style="list-style-type: none"> <li>Plays a role in cell structure in lipids in the cell membranes</li> <li>Promotes brain and memory functions</li> <li>Gives to own manufacture in the body</li> </ul>	Egg yolks, milk, peanuts, soy, wheat germ, livers (beef, veal and turkey)	When low during pregnancy, an increased risk of birth defects; low choline leads to increased risk of cardiovascular disease
	9–18 years	375–550	375–400	2,000–3,000			
	19–50 years	550	425	3,500			
	51–70 years	550	425	3,500			
<b>Vitamin B<sub>12</sub></b> (µg/d) (micrograms*/day) <i>Cobalamin</i>	1–8 years	<b>0.9–1.2</b>	<b>0.9–1.2</b>	None determined	<ul style="list-style-type: none"> <li>Helps maintain nerve tissue and normal blood formation</li> <li>Regeneration of folate</li> </ul>	Animal foods: organ meats, muscle meats, fish, poultry, eggs, milk; fortified cereals	Anemia, neurologic disorders
	9–18 years	<b>1.8–2.4</b>	<b>1.8–2.4</b>				
	19–50 years	<b>2.4</b>	<b>2.4</b>				
	51–70 years	<b>2.4</b>	<b>2.4</b>				

<b>Folate</b> (µg/d)	1–8 years	<b>150–200</b>	<b>150–200</b>	300–400	• Helps red blood cells mature	Organ meats, deep green leafy vegetables,	Anemia, fatigue, gastrointestinal disturbances,
<i>Folic acid</i>	9–18 years	<b>300–400</b>	<b>300–400</b>	600–800	• Interrelated with vitamin B <sub>12</sub> utilization	muscle meats, poultry, fish, eggs, whole-	inadequate intake during pregnancy related to
<i>Folicin</i>	19–50 years	<b>400</b>	<b>400</b>	1,000	• Folic acid supplement*** during pregnancy	grain or fortified cereals	neural tube birth defects (NTD)*
	51–70 years	<b>400</b>	<b>400</b>	1,000	recommended		
<b>Biotin</b> (µg/d)	1–8 years	8–12	8–12	None determined	• Coenzyme* in synthesis of fat, glycogen	Liver, and smaller amounts in meats and	Because data on biotin's adverse effects are
	9–18 years	20–25	20–25		(carbohydrate stored in muscle and liver), and	fruits	limited, caution may be needed
	19–50 years	30	30		amino acids (protein building blocks)		
	51–70 years	30	30				
<b>Fat-soluble vitamins</b>		<b>RDA*/AI*</b>		<b>UL*</b>	<b>Functions in the body</b>	<b>Sources</b>	<b>Deficiency</b>
		<b>Males</b>	<b>Females</b>	<b>M + F</b>			
<b>Vitamin A</b> (µg/d RAE*)	1–8 years	<b>300–400</b>	<b>300–400</b>	600–900	• Promotes growth and normal vision, and protects	Dark leafy green or deep yellow vegetables	Faulty bone and tooth development in infants,
<i>Retinol, Retinal Carotene</i>	9–18 years	<b>600–900</b>	<b>600–700</b>	1,700–2,800	against night blindness	(carrots, winter squash, cushaw, pumpkin,	poor growth, xerophthalmia*, night blindness
*Retinol Activity	19–50 years	<b>900</b>	<b>700</b>	3,000	• Helps keep skin and mucous membrane linings	sweet potatoes); yellow fruits (peaches,	
Equivalent:	51–70 years	<b>900</b>	<b>700</b>	3,000	healthy and resistant to infection	cantaloupe, apricots); liver, fish liver oils,	
1 RAE = 1 µg Retinol					• Large amounts are toxic	dairy foods, butter, margarine, egg yolks	
<b>Vitamin D</b> (iu/d)	1–8 years	600	600	4,000	• Synthesized in skin by ultraviolet light	Fish liver oils and flesh, fortified milk,	Rickets* (soft, fragile bones, enlarged joints,
<i>D Calciferol</i>	9–18 years	600	600	4,000	• Functions to regulate amount of calcium/	exposure to sunlight. Minute amounts in	bowed legs); chest, spinal and pelvic bone
<i>D<sub>2</sub> Ergocalciferol</i>	19–50 years	600	600	4,000	phosphorus absorbed in the blood to mobilize	butter, liver, egg yolk, salmon and sardines	deformities; convulsions; osteomalacia*
<i>D<sub>3</sub> Cholecalciferol</i>	51–70 years	600	600	4,000	and mineralize the bone		
					• Large amounts are toxic		
					• Needed to fight off bacteria and viruses		
<b>Vitamin E</b> (mg/d)	1–8 years	<b>6–7</b>	<b>6–7</b>	200–300	• Not stored in body to any extent	Plant tissues: wheat or rice germ, vegetable	Anemia in premature infants, problems of
<i>Alpha<sup>2</sup>-, beta-,</i>	9–18 years	<b>11–15</b>	<b>11–15</b>	600–800	• Related to action of selenium	oils, green leafy vegetables, nuts, legumes;	nervous system
<i>gamma-tocopherol</i>	19–50 years	<b>15</b>	<b>15</b>	1,000	• Reduces oxidation of vitamin A, carotenes and	meats (other animal foods are poor sources)	
	51–70 years	<b>15</b>	<b>15</b>	1,000	polyunsaturated fatty acids		
<b>Vitamin K</b> (µg/d)	1–8 years	30–55	30–55	None determined	• Bile is necessary for absorption of the vitamin	Deep green leaves (alfalfa, spinach,	Prolonged clotting time, hemorrhagic* disease in
<i>Phylloquinone (K<sub>1</sub>)</i>	9–18 years	60–75	60–75		• Needed to form prothrombin in blood	cabbage), liver, egg yolk, butterfat, (is	newborn infants
<i>Menaquinone (MK<sub>n</sub>)</i>	19–50 years	120	90		• Sulfa drugs and antibiotics interfere with	synthesized in intestine by beneficial	
<i>Menadione</i>	51–70 years	120	90		absorption	bacteria)	
					• Large amounts are toxic		
<b>Minerals/Elements</b>		<b>RDA*/AI*</b>		<b>UL*</b>	<b>Functions in the body</b>	<b>Sources</b>	<b>Deficiency</b>
		<b>Males</b>	<b>Females</b>	<b>M + F</b>			
<b>Calcium</b> (mg/d)	1–8 years	700–1,000	700–1,000	2,500	• Needed to build bones and teeth; helps clot blood	Milk, cheese, ice cream, greens (kale,	Retarded bone mineralization, fragile bones,
	9–18 years	1,300	1,300	2,500	• Helps muscles contract and relax normally. Delays	broccoli, collards, turnips, mustard), dried	rickets*, osteomalacia*, osteoporosis*
	19–50 years	1,000	1,000	2,500	fatigue	peas and beans, fortified juice, soy milk	
	51–70 years	1,200	1,200	2,500			
<b>Chromium</b> (µg/d)	1–8 years	11–15	11–15	None determined	• Works along with insulin in carbohydrate, protein	Brewer's yeast, liver, meat, cheese, whole-	Inability of cells to use glucose for energy
	9–18 years	25–35	21–24		and fat metabolism; glucose tolerance factor	grain cereals, broccoli	
	19–50 years	35	25		(GTF)*		
	51–70 years	30	20				
<b>Copper</b> (µg/d)	1–8 years	<b>340–440</b>	<b>340–440</b>	1,000–3,000	• Aids absorption and use of iron to form	Liver, shellfish, meats, nuts, legumes,	Anemia
	9–18 years	<b>700–890</b>	<b>700–890</b>	5,000–8,000	hemoglobin in red bloods cells	whole-grain cereals	
	19–50 years	<b>900</b>	<b>900</b>	10,000			
	51–70 years	<b>900</b>	<b>900</b>	10,000			
<b>Fluoride</b> (mg/d)	1–8 years	0.7–1	0.7–1	1.3–2.2	• Makes teeth resistant to decay; most effective in	Water (1 part per million is added to some	None known
	9–18 years	2–3	2–3	10	young children	municipal water supplies)	
	19–50 years	4	3	10	• Moderate levels in bone may reduce		
	51–70 years	4	3	10	osteoporosis*		
<b>Iodine</b> (µg/d)	1–8 years	<b>90</b>	<b>90</b>	200–300	• Integral part of thyroid hormones: thyroxine and	Iodized table salt (76 µg/g of salt), seafood,	Cretinism (stunted growth with mental
	9–18 years	<b>120–150</b>	<b>120–150</b>	600–900	triiodothyronine	plants grown in iodine-rich soils, dairy	retardation); endemic goiter
	19–50 years	<b>150</b>	<b>150</b>	1,100		products	
	51–70 years	<b>150</b>	<b>150</b>	1,100			
<b>Iron</b> (mg/d)	1–8 years	<b>7–10</b>	<b>7–10</b>	40	• Part of blood hemoglobin and myoglobin	Liver, organ meats, meat, poultry, egg yolk,	Anemia (frequent in infants, preschool children,
	9–18 years	<b>8–11</b>	<b>8–15</b>	45	• Enzyme involved in energy metabolism	enriched and whole-grain breads, cereals,	teenage girls and pregnant women)
	19–50 years	<b>8</b>	<b>18</b>	45	• Involved in oxygen transport	legumes, dark green vegetables, black strap	
	50–70 years	<b>8</b>	<b>8</b>	45		molasses, peaches, apricots, raisins, prunes,	
						oysters	

<b>Magnesium</b> (mg/d)	1–8 years 9–18 years 19–50 years 51–70 years	<b>80–130</b> <b>240–410</b> <b>400–420</b> <b>420</b>	<b>80–130</b> <b>240–360</b> <b>310–320</b> <b>320</b>	65–100 350 350 350	<ul style="list-style-type: none"> <li>• Activates enzymes involved in protein synthesis</li> <li>• Helps muscles and nerves work</li> <li>• Helps regulate blood sugar levels and promotes normal blood pressure</li> </ul>	Whole-grain cereals, nuts, legumes, meats, milk, green leafy vegetables	Tremors, growth failure
<b>Manganese</b> (mg/d)	1–8 years 9–18 years 19–50 years 51–70 years	1.2–1.5 1.9–2.2 2.3 2.3	1.2–1.5 1.6 1.8 1.8	2–3 6–9 11 11	<ul style="list-style-type: none"> <li>• Activates many enzymes used in carbohydrate and protein metabolism</li> <li>• Bone formation</li> </ul>	Legumes, whole-grain cereals, nuts, tea	None known
<b>Phosphorus</b> (mg/d)	1–8 years 9–18 years 19–50 years 51–70 years	<b>460–500</b> <b>1,250</b> <b>700</b> <b>700</b>	<b>460–500</b> <b>1,250</b> <b>700</b> <b>700</b>	3,000 4,000 4,000 4,000	<ul style="list-style-type: none"> <li>• Builds strong bones and teeth</li> <li>• Releases energy from fat, protein and carbohydrates during metabolism</li> <li>• Aids in formation of genetic material, cell membranes and enzymes</li> </ul>	Breads, cereals, lima beans, meat, poultry, fish, meat alternates, milk, cheese, yogurt	Found widely in foods, so deficiency is rare. Bone loss characterized by weakness, anorexia*, malaise, and pain
<b>Selenium</b> (µg/d)	1–8 years 9–18 years 19–50 years 51–70 years	<b>20–30</b> <b>40–55</b> <b>55</b> <b>55</b>	<b>20–30</b> <b>40–55</b> <b>55</b> <b>55</b>	90–150 280–400 400 400	<ul style="list-style-type: none"> <li>• Antioxidant</li> <li>• Lessens breakdown of vitamin E</li> </ul>	Organ meats, seafoods, cereal foods and plants grown in selenium-rich soil	Hair and nail brittleness and loss
<b>Zinc</b> (mg/d)	1–8 years 9–18 years 19–50 years 51–70 years	<b>3–5</b> <b>8–11</b> <b>11</b> <b>11</b>	<b>3–5</b> <b>8–9</b> <b>8</b> <b>8</b>	7–12 23–34 40 40	<ul style="list-style-type: none"> <li>• Component of many enzymes (carbonic anhydrase and anhydrase carboxypeptidase) and proteins</li> <li>• Controls information from gene to gene so living things develop and function</li> <li>• Plays role in immune function, protein synthesis, and wound healing.</li> </ul>	Seafoods, liver and other organ meats, meats, fish, wheat, yeast. Plant foods are generally low in zinc	Poor wound healing, decreased taste ability
<b>Electrolytes</b>		<b>RDA*/AI*</b>		<b>UL*</b>	<b>Functions in the body</b>	<b>Sources</b>	<b>Deficiency</b>
		<b>Males</b>	<b>Females</b>	<b>M + F</b>			
<b>Sodium</b> (g/d) <sup>4</sup>	1–8 years 9–18 years 19–50 years 51–70 years	1–1.2 1.5 1.5 1.3	1–1.2 1.5 1.5 1.3	1.5–1.9 2.2–2.3 2.3 2.3	<ul style="list-style-type: none"> <li>• Found in extracellular fluid (blood)</li> <li>• Maintains fluid balance and nerve transmission</li> </ul>	Table salt, cheddar cheese, ham, snack foods, most processed foods, salt (sodium chloride) and sodium benzoate/phosphate are added	Fatigue caused by profuse sweating, vomiting and diarrhea
<b>Chloride</b> (g/d)	1–8 years 9–18 years 19–50 years 51–70 years	1.5–1.9 2.3 2.3 2	1.5–1.9 2.3 2.3 2	2.3–2.9 3.4–3.6 3.6 3.6	<ul style="list-style-type: none"> <li>• Helps maintain normal pH of blood (7.35)</li> <li>• Maintains fluid balance and nerve transmission</li> </ul>	Table salt (sodium chloride), barley, wheat, green leafy vegetables, melon, pineapple	Heat cramps, hair loss, tooth loss
<b>Potassium</b> (g/d) <sup>4</sup>	1–8 years 9–18 years 19–50 years 51–70 years	3–3.8 4.5–4.7 4.7 4.7	3–3.8 4.5–4.7 4.7 4.7	None determined	<ul style="list-style-type: none"> <li>• Found inside the cell</li> <li>• Maintains fluid balance and nerve transmission</li> </ul>	Bananas, orange juice, most fruits, potatoes, dried peas, peanuts, nuts, dairy products, and meats	Weakness, poor muscle tone, heart abnormalities, apathy (lack of energy)
<b>Water</b> (liters/day)	1–8 years 9–18 years 19–50 years 51–70 years	1.3–1.7 2.4–3.3 3.7 3.7	1.3–1.7 2.1–2.3 2.7 2.7	None determined	<ul style="list-style-type: none"> <li>• Transports nutrients</li> <li>• Transports waste</li> <li>• Lubricates joints</li> <li>• Regulates body temperature</li> <li>• Cell hydration</li> </ul>	Water, juices, beverages, high-moisture solid foods (soups, watermelon, meats, etc.)	Dehydration, constipation

\* See Glossary for definitions

\*\*Average minimum amounts of glucose used by brain

\*\*\*Supplement during pregnancy of 400 µg or mcg folic acid plus folate intake of a varied diet

<sup>1</sup> NE (niacin equivalent) is equal to 1 mg of niacin or 60 mg of dietary tryptophan

<sup>2</sup> RAE = Retinol activity equivalents. 1 retinol equivalent = 1 µg retinol or 6 µg beta-carotene

<sup>3</sup> α-tocopherol includes the only form (RRR-α-tocopherol) that occurs naturally in foods and with variations of this form in fortified foods and supplements.

<sup>4</sup> Estimated sodium and potassium minimum requirements. AI\* has been set for healthy individuals and the UL\* may be too high for persons with hypertension.

#### Texas A&M AgriLife Extension Service

*AgriLifeExtension.tamu.edu*

More Extension publications can be found at *AgriLifeBookstore.org*

Educational programs of the Texas A&M AgriLife Extension Service are open to all people without regard to race, color, sex, disability, religion, age, or national origin.

The Texas A&M University System, U.S. Department of Agriculture, and the County Commissioners Courts of Texas Cooperating.

Produced by Texas A&M AgriLife Communications